



PATIENT NAME

PATIENT ID

PATIENT DOB

## Remdesivir and TACTIC -R Trial

Treatment of patients with Remdesivir is not a contraindication to participation in TACTIC-R, indeed its use is encouraged as standard of care, if the patient is eligible.

To check if your patient is eligible for Remdesivir, TACTIC-R, or both:

1. **Must be a hospital in-patient, not receiving invasive mechanical ventilation, ECMO, or cardiovascular support (vasopressors or inotropes)**

### 2. Complete the ELIGIBILITY CRITERIA TICK BOX CHARTS

If all boxes are ticked for Remdesivir, then refer to respiratory to ensure there are no contraindications to treatment.

If all boxes are ticked for TACTIC-R, then refer to the TACTIC-R team to ensure there are no contraindications to treatment.

If all boxes are ticked for both Remdesivir and TACTIC-R, then your patient can be considered for **both**.

\_\_\_\_\_ Sign and Date



<b>PATIENT NAME</b>
<b>PATIENT ID</b>
<b>PATIENT DOB</b>

**Prescreening Checklists for Determining if a Patient is Eligible for Remdesivir and/or TACTIC-R**

<b>Eligibility Criteria for Remdesivir</b>	<b>TICK if YES</b>
Age over 12 years	
Confirmed COVID with positive PCR result within the preceding 72 hours	
Weight ≥40kg	
Creatinine clearance ≥50 ml/min	
ALT <250 U/L	
No evidence of severe chronic liver disease (Child Pugh score 10 or more. You will need bilirubin, albumin, INR, presence ascites and encephalopathy: <a href="https://www.mdcalc.com/child-pugh-score-cirrhosis-mortality">https://www.mdcalc.com/child-pugh-score-cirrhosis-mortality</a> )	
Requiring ≥40% oxygen to maintain oxygen saturations ≥94% on two occasions > 1 hour apart OR Within 24h commencing CPAP or HFNO2 to maintain O2 sats >94% and have not been previously mechanically ventilated for Covid19 OR Just starting ventilation, in early phase of infection and potential benefit determined by multidisciplinary assessment	
<b>ONLY if ALL boxes ticked, discuss with respiratory consultant as may be suitable for Remdesivir</b>	

<b>Eligibility Criteria for TACTIC-R Trial</b>	<b>TICK if YES</b>
Age over 18 years	
Clinical picture strongly suggestive of COVID, <b>with or without</b> a positive Covid-19 test	
Not pregnant or breastfeeding	
Not participating in RECOVERY or REMAP-CAP	
Not currently being treated with probenecid, chronic IVIG, Baricitinib or Ravulizumab	
Not known to have active TB, unresolved Neisseria meningitidis infection, active Hepatitis B/C or Varicella Zoster (no blood testing required to exclude these)	
Creatinine clearance ≥30 ml/min	
ALT ≤250 U/L	
VTE prophylaxis or maintenance therapy not contraindicated	
<b>ONLY if all boxes ticked, may be suitable for TACTIC-R, please discuss with TACTIC-R team</b>	